


VYALEV Infusion Site Tracker



Use this tracker to help you note down the location and date of your last infusion site

It may be helpful to place this Infusion Site Tracker on your fridge or in the area of your home where you administer your VYALEV treatment. You can download more copies of the tracker from vyalev.com.au using the access code: **connect** or by scanning the **QR code**.



The 6 sites in the diagram above are for illustration purposes and serve as a guide only. You are not restricted to using these infusion sites; this is based on changing your infusion site every 3 days.

Your infusion site should be at least 5 cm from any areas of scarred or hardened tissue, stretch marks, skin folds or creases where the body naturally bends (e.g. while sitting or exercising), or areas where clothing might cause irritation (e.g. near the belt line).

Turn to the back page for important steps and reminders about rotating your infusion site.





Important steps for rotating your infusion site

Step 1:

Before changing your infusion site, wash your hands with soap and water then dry them with a paper towel.

Step 2:

Carefully remove the old infusion set (tubing, insertion device and cannula). Dispose of the infusion set in the disposable sharps container.

Step 3:


Choose a new infusion site that is **at least 2.5 cm** away from a site used in the last 12 days and **at least 5 cm** from your belly button.

Step 4:

Prepare the skin around the new infusion site with an alcohol pad before inserting the new cannula.

Helpful tips and reminders


If needed, remove hair around the infusion site to ensure proper attachment of the adhesive tape on the cannula to your skin.



You may need to adjust your belt line so that it falls below the infusion sites.



You may need to rotate your infusion site more often than every 3 days, if you notice any of the skin changes mentioned below or if advised by your prescribing specialist.



Call your prescribing specialist straight away if you notice any skin changes at the infusion site, such as redness, warmth, swelling, pain, and discolouration when you apply pressure to it.

For more information about your VYALEV medicine and its delivery system, speak with your prescribing specialist and refer to the VYALEV Instructions for Use and Consumer Medicine Information, available at abbv.ie/au-vylv-cmi, by scanning the QR code or from your pharmacist.

