

# VYALEV Step-by-step Infusion Guide

Your prescribing specialist will have gone through how to set up and administer VYALEV. Use this resource as a guide to help organise your infusion supplies and follow the correct steps when administering your medicine. Review the Consumer Medicine Information and Instructions for Use provided. If you have any questions, contact your prescribing specialist.



## 1. Wash your hands

Always wash your hands with soap and water and dry them with a paper towel. Also, make sure the surface you are working on is clean. This will help to avoid contamination.

## 2. Gather your supplies

Gather VYALEV and supporting items as listed below and place them on a clean, flat surface such as a tray or table.



Solution vial



Insertion device  
and cannula



Vial adapter



Syringe




Tubing



VYAFUSER® Pump



2 Alcohol pads



Paper towels\*



Disposable sharps  
container\*

Once you have gathered all the items needed for your treatment, flip the page down for the administration steps.

**Inspect the items for damage and check the expiration date as you gather your supplies.**

- ✗ Do not use any components if their sterile packaging has been damaged prior to use.
- ✗ Do not use the VYALEV solution, vial adapter or syringe if it is expired.

\*Items not provided.

# VYALEV Step-by-step Infusion Guide

Your prescribing specialist will have gone through how to set up and administer VYALEV. Use this resource as a guide to help organise your infusion supplies and follow the correct steps when administering your medicine. Review the Consumer Medicine Information and Instructions for Use provided. If you have any questions, contact your prescribing specialist.



## 3. Prepare the solution vial




- Remove the solution vial from the refrigerator and allow to sit at room temperature out of direct sunlight for **30 minutes**.



- Do not** warm the solution vial in any way other than letting it warm at room temperature. For example, do not warm in microwave or in hot water.
- Do not** use if the solution is cloudy or contains flakes or particles.
- Once you have inspected the solution, remove the solution vial cap. Wipe the top of the vial stopper with an alcohol pad and allow to dry. This will help to avoid contamination.

## 4. Prepare the syringe



- Take a new syringe and remove it from its packaging.
- Depress the rubber plunger to fully expel all air.

## 5. Attach the vial adapter to the solution vial



- Remove the peel-off cover from the packaging containing the sterile vial adapter.
- If the tip of the vial adapter comes into contact with an unclean surface, discard it and get a new one.




- Keeping the vial adapter in the packaging, push the adapter down onto the top of the solution vial until you hear a “snap”.




- Once the adapter is attached to the vial, remove the protective cap.

**If the tip of the vial adapter or syringe comes into contact with an unclean surface, discard it and get a new one.**


## 6. Transfer medicine from the vial into the syringe




Attach the syringe to the vial adapter by pushing and then screwing it into place. **Do not** overtighten.



Hold the syringe vertically with the solution vial directly above.



Transfer the solution from the vial into the syringe by pulling down the plunger. Always withdraw the entire contents of the solution vial into the syringe (around the 12 mL mark), or until you see air at the tip of the syringe.




Remove large air bubbles from the syringe. The presence of air may affect how accurately the dose is delivered.

The syringe may be unscrewed from the vial adapter once large air bubbles have been removed.

Please refer to the VYALEV Instructions for Use for detailed steps on how to remove air bubbles from the syringe.


Diagrams are for illustration purposes and your vial adapter may look different.

## 7. Connect your tubing to the filled syringe




- Place the syringe on a clean surface, making sure the syringe tip does not come in contact with an unclean surface.
- Remove the infusion set tubing from the package.
- Connect the infusion set tubing to the syringe and lay the infusion site connector end of infusion set tubing on a clean paper towel before preparing to prime the infusion line.

## 8. Prepare the VYAFUSER® Pump



- Turn on pump display and place the syringe inside the pump. Then, close the pump and invert it.
- Follow the on-screen instructions to prime the infusion set.
- Once a drop of VYALEV solution appears on the connector site, the infusion set tubing has been primed.
- It is important to wait for one minute to ensure the solution has stopped dripping from the connector site and is free from drops before attaching it to the cannula. Otherwise, it may later be difficult to detach from the cannula.


## 9. Insert the cannula into the body



- Select your desired infusion site – at least 5 cm from the belly button and at least 2.5 cm from the previous insertion site.
- Wipe the infusion site with an alcohol pad and allow to dry for a minute.
- Insert the cannula into the skin.


**Change the infusion site every time you change the infusion set, trying not to repeat the last used site for at least 12 days.**

## 10. Connect the infusion set tubing to the cannula



Attach the connector site of the infusion set to the cannula. Make sure it snaps securely in place to prevent leakage.

## 11. Disposal



After you have administered VYALEV, place your used vial, tubing and cannula into the disposal sharps container straight away. Similarly, when replacing the syringe, discard the used syringe in the disposable sharps container.

**This resource is for demonstration and training purposes only. For more detailed instructions, refer to the appropriate Instructions for Use manuals (IFUs), Consumer Medicine Information (CMI), Administration Guide, or speak with your prescribing specialist. Please ensure you have read the IFUs and CMI before using VYALEV.**



**VYALEV<sup>®</sup>**  
**foslevodopa/foscarbidopa**  
Solution for subcutaneous infusion  
240 mg/12 mg per mL